

**GENETIC DIVERSITY AND SUSCEPTIBILITY OF THE SELECTED
VISAYAS POPULATIONS OF BROWN PLANTHOPPER,
Nilaparvata lugens (Stål) (HEMIPTERA: DELPHACIDAE)
TO *Beauveria bassiana* (Bals.-Criv.) Vuill.**

Jenuelle Kaye P. Arabit¹, Melissa P. Montecalvo², and Gelyn D. Sapin^{1*}

¹Institute of Weed Science, Entomology and Plant Pathology, College of Agriculture and Food Science, University of the Philippines Los Baños, College, Laguna, Philippines

²National Crop Protection Center, College of Agriculture and Food Science, University of the Philippines Los Baños, College, Laguna, Philippines

*Corresponding author: gdsapin@up.edu.ph

ABSTRACT

The brown planthopper (BPH), *Nilaparvata lugens* (Stål), is a notorious insect pest of rice due to its ability to transmit viruses such as rice grassy stunt virus and rice ragged stunt virus. Its direct feeding in rice can cause hopperburn, which is described as wilting of the crop, causing a decline in yield. Investigating the genetic diversity of BPH populations in the Philippines is crucial for improving pest control strategies. Hence, this study aims to determine the genetic diversity of the different BPH populations collected from various provinces in the Visayas region of the Philippines and determine their susceptibility to an entomopathogenic fungus, *Beauveria bassiana* (Bals.-Criv.) Vuill. The study was conducted from December 2022 to December 2023 using the BPH collected from different localities in Cebu, Bohol, and Negros Oriental. A total of fourteen sequences amplifying the *cytochrome C oxidase I* gene were analyzed. The nucleotide and haplotype diversities of all the sequences analyzed were 0.00276 and 0.780, respectively. Based on the molecular analysis, five distinct haplotypes were observed with moderate genetic differentiation ($F_{st} = 0.08680$). Analyses of molecular variance showed that 91.32% of variation was due to the difference within populations. The different populations of BPH exhibited similar degrees of susceptibility to *B. bassiana* at 1×10^8 conidia/mL with up to 60% adult mortality and an LT50 of 4.08-4.26 days under laboratory conditions. The result may suggest that the little to moderate genetic difference among the BPH populations used in the study does not significantly affect their susceptibility to *B. bassiana*.

Keywords: *Nilaparvata lugens*, genetic diversity, *cytochrome c oxidase I*, entomopathogenic fungi, *Beauveria bassiana*

INTRODUCTION

Rice (*Oryza sativa* L.) is considered a staple food that provides more than 20% of the calories consumed worldwide (Sharif et al., 2014). Globally, the Philippines ranks eight among the world's top rice producers and sixth among the world's top rice consumers (USDA, 2023). Due to insufficient local production (Mamiit et al., 2021), the Philippines is currently the largest global importer of rice (USDA, 2023). One of the major constraints in rice production is insect pests because they can cause yield losses (Mondal et al., 2017). Brown planthopper (BPH), *Nilaparvata lugens* (Stål) (Hemiptera: Delphacidae), is one of the most important key insect pests of rice (Sanchez, et al., 1980; Ghaffar et al., 2011; Hereward et al., 2020). This insect transmits viral diseases such as rice grassy stunt virus and rice ragged stunt virus (Cabauatan et al., 2009). Its direct feeding causes hopperburn, which often resembles senescence that includes wilting, chlorosis, and stunted growth of the crop (Backus et al., 2005). Annually, its damage has caused an estimated loss of more than \$300 million in rice production losses in Asia (Min et al., 2014).

Insecticide application is the most widely used and most effective control for managing BPH (Rahman & Jahan, 2006). However, misuse and overuse of these products have caused several drawbacks due to the ability of the BPH to develop insecticide resistance (Heong, 2009; Khoa et al., 2019). Furthermore, excessive use of insecticides would coincidentally kill the natural enemies, thereby reducing the presence of natural control (Bottrell & Schoenly, 2012). Due to the disadvantages of chemical control, the use of biological control has been regarded as one of the most promising technologies for sustainable agriculture (Tracy, 2014). Entomopathogenic fungi (EPF) are biological control agents that infect and often kill insects and other arthropods through cuticle penetration (Mantzoukas et al., 2022). *Beauveria bassiana* (Bals.-Criv.) Vuill. is a widely known species of EPF that is currently being utilized as a biological control agent against several arthropods (de Faria & Wraight, 2007). It is distributed worldwide (Zimmermann, 2007) with an extensive host range (Ownley et al., 2004) and can control insect pests without harming non-target organisms (Skinner et al., 2014; Mascarin & Jaronski, 2016; Dannon et al., 2020).

BPH populations vary in their physiological and ecological characteristics (Nagata & Matsuda, 1980). The genetic differences between BPH populations were observed by Mun et al. (1999). Identifying the differences among local BPH populations can be analyzed using mitochondrial genome sequences

(Matsumoto et al., 2013), which has become a widely used molecular marker to further understand the genetic structure (Roderick, 1996) and phylogenetic relationships in insects (Simon et al., 1994). One of the well-known mitochondrial genes, the gene encoding for the component I of *cytochrome oxidase (COI)* (Mun et al., 1999), has additional properties that make it particularly suited as a molecular marker for evolutionary studies (Lunt et al., 1996). *COI* is frequently used as a molecular marker since no other gene can be found in several taxonomic groups (Deagle et al., 2014). Besides its universality, this gene region was shown to be highly conserved within species while also being sufficiently variable between species, allowing reliable species identification (Pentinsaari et al., 2016).

Laquinta et al. (2019) have preliminary results in understanding the genetic structure of BPH in the different localities in Luzon. They used the *COI* gene region and reported the occurrence of polymorphism and numerous haplotypes in the Luzon population. In the study of Mun et al. (1999), the *COI* region of the BPH individuals collected from 11 localities in Asian countries was analyzed and reported the presence of three distinct haplotypes, and these haplotypes were all present in the sampled populations from the Philippines. Mashhoor et al. (2015) used the *COI* region to determine the genetic structure and phylogenetic status of BPH in Kerala, India. They reported similarities between BPH isolated in Kerala, India, with the population isolated from Andhra Pradesh and some populations isolated from Japan and China. In Indonesia, Winnie et al. (2020) combined the *COI* and *COII* regions to analyze six populations of BPH isolated from Java, Indonesia, where it showed that the populations have high haplotype diversity but low nucleotide diversity.

The ability of the insect to adapt to their local environment, resulting in the occurrence of variations within and among local populations, could limit the effectiveness of the current pest management practice against BPH. Hence, understanding the genetic diversity of the BPH population shall help design more efficient pest management practices against BPH. Here, we analyze the genetic diversity of the selected Visayas BPH populations and determine the susceptibility of these BPH populations to *B. bassiana*.

MATERIALS AND METHODS

Insect Samples

BPH samples were collected from June to July 2022 from the different fields located in the three provinces in Visayas, Philippines, namely, Asturias, Cebu (10.55546 °N, 123.72809 °E); Sierra Bullones, Bohol (coordinates not recorded); and Caticugan, Siaton, Negros Oriental (9.07914 °N, 123.02309 °E).

Amplification of *COI* Gene of Brown Planthopper

The DNA of an individual BPH adult was isolated following the manufacturer's protocol for the Animal Tissue Genomic DNA Purification Kit (Promega Wizard®, Madison, Wisconsin, USA) with some modifications, wherein the Proteinase K was incubated upon the addition of RNase for 30 mins at 37°C; cold isopropanol was used; and all centrifugation steps during DNA precipitation and rehydration were increased to 10 mins. The quantity and quality of the isolated DNA samples were determined using a UV/VIS spectrophotometer (BioDrop™). The *cytochrome c oxidase subunit I (COI)* gene amplification was done with the use of the following set of primers: CI-J-2183 5'-CAA CAT TTA TTT TGA TTT TTT GG-3' and TL2-N-3014 5'-TCC AAT GCA CTA ATC TGC CAT ATT A-3' (Mun et al., 1999). The polymerase chain reaction was carried out using a 25 µL PCR reaction mixture, which contained 12.5 µL of 2x Taq Master Mix (Vivantis, www.vivantistechnologies.com), 1 µL each of forward and reverse primers with a concentration of 10 pmol/µL, 8.5 µL of nuclease-free water, and 2 µL of 50 ng DNA template. The thermal profile used for amplification was 35 cycles of 94°C for 1 min, a denaturation step of 48°C for 30 s, and an annealing step of 72°C for 45 sec (Mun et al., 1999). The amplified *COI* gene fragments were resolved in 1.5% agarose gel, stained with SYBR Safe DNA gel stain (Invitrogen, Carlsbad, California, USA), and the molecular size of the amplicons was estimated using the VC 100bp Plus DNA ladder (Vivantis, www.vivantistechnologies.com).

COI Gene Nucleotide Sequence Analyses

The amplified DNA samples were sent to Macrogen, Inc. (Seoul, South Korea) through the sequencing services of Kinovett Scientific Solutions Co. (Quezon City, Philippines). The obtained nucleotide sequences were aligned together with the *COI* gene sequence of *N. lugens* obtained from GenBank (Accession Number: NC_021748.1) using Clustal W (Thompson et al., 1994). The sequences were edited and trimmed into 698 base pairs using BioEdit Sequence

Alignment Editor Version 7 (Hall, 1999). The translation of the nucleotide sequences corresponding to 227 amino acid residues was carried out using ExPASy (<https://web.expasy.org/translate/>) (Gasteiger et al., 2003). The genetic diversity parameters, including the number of polymorphic sites, the number of haplotypes, transitions, transversion, nucleotide diversity, haplotype diversity, and nucleotide composition were estimated using DnaSP Version 6 (Rozas et al., 2017).

The pairwise p distances, pairwise F_{st} , and analyses of molecular variance (AMOVA) were determined using Arlequin Version 3 (Excoffier et al., 2005). The COI haplotypes of *N. lugens* were compared to the existing COI sequences of *N. lugens* in GenBank using the BLASTn program (<http://blast.ncbi.nlm.nih.gov>). The TCS haplotype networks were constructed using the TCS Network (Clement et al., 2000) to examine the relationships between the haplotypes.

Rearing of Brown Planthopper and Mass Production of *Beauveria bassiana*

The BPH populations were reared separately in cages at the National Crop Protection Center, College of Agriculture and Food Science, University of the Philippines Los Baños (NCPC-CAFS, UPLB), College, Laguna, Philippines. The BPH nymphs and adults collected from the three provinces in Visayas, Philippines, were fed with a susceptible variety of rice seedlings, Taichung Native 1 (TN1), grown in pots supplied in each cage, wherein 30-day-old rice seedlings planted in pots are placed inside a wire mesh cage. Adult BPH was used for biological assays of the entomopathogenic fungus. The entomopathogenic fungus, *B. bassiana*, was sourced from the Mycology Laboratory, NCPC-CAFS, UPLB, which was originally isolated from rice bug (*Leptocorisa oratorius* (Fabricius)). The fungus was subcultured in Potato Dextrose Agar (PDA) until pure cultures were obtained.

Mortality Assay against *N. lugens*

The conidial suspension with a concentration of 1×10^8 conidia/mL was prepared by diluting the stock solution in 0.1% Tween 80 solution. The three BPH populations were exposed to the conidial suspension of *B. bassiana*. Rice seedlings were washed in running water and sterilized in rinses of 0.5% sodium hypochlorite and sterile distilled water. After which, *B. bassiana* suspension and 0.1% Tween 80 solution were sprayed on rice seedlings in fungal-treated and control set-ups, respectively. Set-ups were placed inside sterile test tubes. Ten adult BPH were introduced to each test tube, and each test tube represents one replicate. A total of six replicates were done in each BPH population. Surface-sterilized fresh rice leaves were introduced after 48 hours.

Data Analyses

The mortality of the BPH was recorded daily. Following Lacey and Brooks (1997), the cadavers were sterilized using 0.5% sodium hypochlorite for 30 seconds and washed twice in sterile distilled water. After that, the cadavers were placed in sterilized Petri dishes with moistened filter paper and stored at room temperature.

The mortality of the BPH was corrected using Abbott's Formula (Abbott, 1925). The data were checked for normality using Shapiro-Wilk test, upon confirming normality a one-way analysis of variance (ANOVA) was done for the percent mean cumulative mortality of the different BPH populations. All the statistical analyses were done using GraphPad Prism (Version 10.1.1).

RESULTS AND DISCUSSION

Genetic diversity analysis of *N. lugens* populations

The *COI* gene regions were successfully amplified from the three populations of BPH collected from Visayas, Philippines, namely, Asturias, Cebu; Sierra Bullones, Bohol; and Caticugan, Siaton, Negros Oriental (Figure 1). The amplicons were approximately 900 bp, which agrees with the reported *COI* gene region of BPH by Mun et al. (1999).




Figure 1. DNA fragments amplified from brown planthopper, *Nilaparvata lugens* (Stål), containing *cytochrome c oxidase I (COI)* gene sequence resolved in 1.5% agarose gel stained with SYBR® Safe DNA gel stain (Invitrogen, Thermo Fisher Scientific, Carlsbad, California, USA). The molecular weights of the PCR products were estimated using a 100 bp plus ladder (Vivantis www.vivantis.com). The DNA templates used were isolated from BPH collected from the Visayas provinces, namely Bohol, Cebu, and Negros Oriental.

The genetic diversity of BPH was analyzed based on the nucleotide sequences of the *COI* gene. Five polymorphic sites were identified at positions 103, 301, 436, 664, and 671, all of which were parsimony-informative sites (Figure 2). Polymorphisms at positions 301 and 671 were translations, while polymorphisms at positions 103, 436, and 664 were transversions. Upon translation using ExPASy, the variations were found to produce silent mutations at amino acids 35, 101, and 146 and missense mutations at amino acids 221 and 223 (Figure 3). The comparison of the diversity indices of the BPH populations is shown in Table 1. The nucleotide and haplotype diversities of all *COI* gene sequences analyzed were 0.00276 ± 0.00047 and 0.780 ± 0.085 , respectively. The haplotype diversities of the different populations had a range of 0.500 to 0.800 (mean = 0.633), whereas the nucleotide diversities ranged from 0.00143 to 0.00430 (mean = 0.00248). Notably, haplotype diversity was relatively higher than nucleotide diversity, suggesting small differences between BPH haplotypes. This observation is consistent with recent population expansion from a low effective population size (Guru-Pirasanna-Pandi et al., 2022).

Table 1. Comparison of the molecular diversity indices of the brown planthopper, *Nilaparvata lugens* (Stål), collected from rice fields in three provinces (Bohol, Cebu, and Negros Oriental) in Visayas, Philippines.

INDICES	COLLECTION SITES			
	BOHOL	CEBU	NEGROS ORIENTAL	ALL
No. of Samples	5	4	5	14
Molecular diversity				
• No. of Polymorphic	2	2	5	5
• No. of Transitions	1	1	2	2
• No. of Transversions	1	1	3	3
• No. of Substitutions	2	2	5	5
Haplotype information (number of samples)	Haplotype 1 (3) Haplotype 2 (2)	Haplotype 1 (3) Haplotype 3 (1)	Haplotype 3 (2) Haplotype 4 (1) Haplotype 5 (2)	
Haplotype Diversity	0.600 ± 0.175	0.500 ± 0.265	0.800 ± 0.164	0.780 ± 0.085
Nucleotide Diversity	0.00172 ± 0.00050	0.00143 ± 0.00076	0.00430 ± 0.00097	0.00276 ± 0.00047
Nucleotide Composition				
• C	16.05	16.01	15.99	16.03
• T	37.45	37.39	37.42	37.43
• A	32.84	32.95	32.87	32.84
• G	13.67	13.65	13.72	13.70

Haplotype-1	AATTTCTCAT	ATTATTATAC	AAGAAAGAGG	AAAAAGGGAA	ACTTTCGGAT	CTATTGGAAT	60
Haplotype-2	AATTTCTCAT	ATTATTATAC	AAGAAAGAGG	AAAAAGGGAA	ACTTTCGGAT	CTATTGGAAT	60
Haplotype-3	AATTTCTCAT	ATTATTATAC	AAGAAAGAGG	AAAAAGGGAA	ACTTTCGGAT	CTATTGGAAT	60
Haplotype-4	AATTTCTCAT	ATTATTATAC	AAGAAAGAGG	AAAAAGGGAA	ACTTTCGGAT	CTATTGGAAT	60
Haplotype-5	AATTTCTCAT	ATTATTATAC	AAGAAAGAGG	AAAAAGGGAA	ACTTTCGGAT	CTATTGGAAT	60
Haplotype-1	AATTTATGCA	ATAATTGCAA	TTGGAAATTT	AGGTTTTATT	GTITGAGCTC	ACCATATATT	120
Haplotype-2	AATTTATGCA	ATAATTGCAA	TTGGAAATTT	AGGTTTTATT	GTITGAGCTC	ACCATATATT	120
Haplotype-3	AATTTATGCA	ATAATTGCAA	TTGGAAATTT	AGGTTTTATT	GTITGAGCTC	ACCATATATT	120
Haplotype-4	AATTTATGCA	ATAATTGCAA	TTGGAAATTT	AGGTTTTATT	GTITGAGCTC	ACCATATATT	120
Haplotype-5	AATTTATGCA	ATAATTGCAA	TTGGAAATTT	AGGTTTTATT	GTITGAGCTC	ACCATATATT	120
Haplotype-1	TACTGTAGGT	ATAGATATTG	ATACCCGAGC	CTATTTTACG	TCAGCTACTA	TAATTATTGC	180
Haplotype-2	TACTGTAGGT	ATAGATATTG	ATACCCGAGC	CTATTTTACG	TCAGCTACTA	TAATTATTGC	180
Haplotype-3	TACTGTAGGT	ATAGATATTG	ATACCCGAGC	CTATTTTACG	TCAGCTACTA	TAATTATTGC	180
Haplotype-4	TACTGTAGGT	ATAGATATTG	ATACCCGAGC	CTATTTTACG	TCAGCTACTA	TAATTATTGC	180
Haplotype-5	TACTGTAGGT	ATAGATATTG	ATACCCGAGC	CTATTTTACG	TCAGCTACTA	TAATTATTGC	180
Haplotype-1	GGTCCCCACC	GGAA TC AAAA	TTTTTAGATG	AATCGCAACA	ATTACGGTT	CCAAAATGAA	240
Haplotype-2	GGTCCCCACC	GGAA TC AAAA	TTTTTAGATG	AATCGCAACA	ATTACGGTT	CCAAAATGAA	240
Haplotype-3	GGTCCCCACC	GGAA TC AAAA	TTTTTAGATG	AATCGCAACA	ATTACGGTT	CCAAAATGAA	240
Haplotype-4	GGTCCCCACC	GGAA TC AAAA	TTTTTAGATG	AATCGCAACA	ATTACGGTT	CCAAAATGAA	240
Haplotype-5	GGTCCCCACC	GGAA TC AAAA	TTTTTAGATG	AATCGCAACA	ATTACGGTT	CCAAAATGAA	240
Haplotype-1	CTTTTCCCCC	CAAA TA ATTT	GATCATTAGG	ATTCATTTTA	CTTTTTACTA	TTGGAGGATT	300
Haplotype-2	CTTTTCCCCC	CAAA TA ATTT	GATCATTAGG	ATTCATTTTA	CTTTTTACTA	TTGGAGGATT	300
Haplotype-3	CTTTTCCCCC	CAAA TA ATTT	GATCATTAGG	ATTCATTTTA	CTTTTTACTA	TTGGAGGATT	300
Haplotype-4	CTTTTCCCCC	CAAA TA ATTT	GATCATTAGG	ATTCATTTTA	CTTTTTACTA	TTGGAGGATT	300
Haplotype-5	CTTTTCCCCC	CAAA TA ATTT	GATCATTAGG	ATTCATTTTA	CTTTTTACTA	TTGGAGGATT	300
Haplotype-1	AACAGGTGTA	ATATTATCAA	ATTCCTCAAT	TGATATTATT	CTACATGATA	CCTATTATGT	360
Haplotype-2	AACAGGTGTA	ATATTATCAA	ATTCCTCAAT	TGATATTATT	CTACATGATA	CCTATTATGT	360
Haplotype-3	AACAGGTGTA	ATATTATCAA	ATTCCTCAAT	TGATATTATT	CTACATGATA	CCTATTATGT	360
Haplotype-4	AACAGGTGTA	ATATTATCAA	ATTCCTCAAT	TGATATTATT	CTACATGATA	CCTATTATGT	360
Haplotype-5	AACAGGTGTA	ATATTATCAA	ATTCCTCAAT	TGATATTATT	CTACATGATA	CCTATTATGT	360
Haplotype-1	AGTGGCTCAT	TTTCACTATG	TCCTTTCCAT	GGGAGCAGTA	TTCAACCATA	TCCGTAGATT	420
Haplotype-2	AGTGGCTCAT	TTTCACTATG	TCCTTTCCAT	GGGAGCAGTA	TTCAACCATA	TCCGTAGATT	420
Haplotype-3	AGTGGCTCAT	TTTCACTATG	TCCTTTCCAT	GGGAGCAGTA	TTCAACCATA	TCCGTAGATT	420
Haplotype-4	AGTGGCTCAT	TTTCACTATG	TCCTTTCCAT	GGGAGCAGTA	TTCAACCATA	TCCGTAGATT	420
Haplotype-5	AGTGGCTCAT	TTTCACTATG	TCCTTTCCAT	GGGAGCAGTA	TTCAACCATA	TCCGTAGATT	420
Haplotype-1	TATCCATTGA	TACCCCTTAT	TTACAGGTAG	AAACATAAAT	AATAAATGAC	TAAAAATTCA	480
Haplotype-2	TATCCATTGA	TACCCCTTAT	TTACAGGTAG	AAACATAAAT	AATAAATGAC	TAAAAATTCA	480
Haplotype-3	TATCCATTGA	TACCCCTTAT	TTACAGGTAG	AAACATAAAT	AATAAATGAC	TAAAAATTCA	480
Haplotype-4	TATCCATTGA	TACCCCTTAT	TTACAGGTAG	AAACATAAAT	AATAAATGAC	TAAAAATTCA	480
Haplotype-5	TATCCATTGA	TACCCCTTAT	TTACAGGTAG	AAACATAAAT	AATAAATGAC	TAAAAATTCA	480
Haplotype-1	ATTTTATTC	ATATTTCTAG	GAGTAAATTT	AACATTTTTT	CCCCAACATT	TTTTAGGATT	540
Haplotype-2	ATTTTATTC	ATATTTCTAG	GAGTAAATTT	AACATTTTTT	CCCCAACATT	TTTTAGGATT	540
Haplotype-3	ATTTTATTC	ATATTTCTAG	GAGTAAATTT	AACATTTTTT	CCCCAACATT	TTTTAGGATT	540
Haplotype-4	ATTTTATTC	ATATTTCTAG	GAGTAAATTT	AACATTTTTT	CCCCAACATT	TTTTAGGATT	540
Haplotype-5	ATTTTATTC	ATATTTCTAG	GAGTAAATTT	AACATTTTTT	CCCCAACATT	TTTTAGGATT	540
Haplotype-1	AACTGGTATA	CCACGACGAT	ACTCTGACTA	TCCAGATATA	TACACCCCTGT	GAAACCCTTT	600
Haplotype-2	AACTGGTATA	CCACGACGAT	ACTCTGACTA	TCCAGATATA	TACACCCCTGT	GAAACCCTTT	600
Haplotype-3	AACTGGTATA	CCACGACGAT	ACTCTGACTA	TCCAGATATA	TACACCCCTGT	GAAACCCTTT	600
Haplotype-4	AACTGGTATA	CCACGACGAT	ACTCTGACTA	TCCAGATATA	TACACCCCTGT	GAAACCCTTT	600
Haplotype-5	AACTGGTATA	CCACGACGAT	ACTCTGACTA	TCCAGATATA	TACACCCCTGT	GAAACCCTTT	600
Haplotype-1	TTCTTCTATG	GGTTCATTCA	TTTCCCTAAT	TAGAATTTTA	ATATTAATGT	TTATTATATG	660
Haplotype-2	TTCTTCTATG	GGTTCATTCA	TTTCCCTAAT	TAGAATTTTA	ATATTAATGT	TTATTATATG	660
Haplotype-3	TTCTTCTATG	GGTTCATTCA	TTTCCCTAAT	TAGAATTTTA	ATATTAATGT	TTATTATATG	660
Haplotype-4	TTCTTCTATG	GGTTCATTCA	TTTCCCTAAT	TAGAATTTTA	ATATTAATGT	TTATTATATG	660
Haplotype-5	TTCTTCTATG	GGTTCATTCA	TTTCCCTAAT	TAGAATTTTA	ATATTAATGT	TTATTATATG	660
Haplotype-1	AGAAAGATTA	AGATTTTAAAC	GAAAAATGGT	GTTTAAAA	698		
Haplotype-2	AGAAAGATTA	AGATTTTAAAC	GAAAAATGGT	GTTTAAAA	698		
Haplotype-3	AGAAAGATTA	AGATTTTAAAC	GAAAAATGGT	GTTTAAAA	698		
Haplotype-4	AGAAAGATTA	AGATTTTAAAC	GAAAAATGGT	GTTTAAAA	698		
Haplotype-5	AGAAAGATTA	AGATTTTAAAC	GAAAAATGGT	GTTTAAAA	698		

Figure 2. Sequence alignment of the five *COI* haplotypes of the brown planthopper, *Nilaparvata lugens* (Stål), collected from rice fields in Bohol, Cebu, and Negros Oriental, Visayas, Philippines.

COI haplotypes can exhibit variability in the biological and ecological traits of an insect. This was demonstrated in the study of Camus et al. (2015), wherein the variation in mitochondrial haplotypes affects the sex and life stages of the insect. Furthermore, the mitochondrial genotype was considered to have a

linkage with aging, fecundity, starvation stress, and thermal tolerance. There are also evolutionary studies that documented the consequences that arise from the variation in the mitochondria (Ballard & Melvin, 2010). These variations that cause mutations could occur in every population. The effects of these mutations may be classified into three types, namely deleterious, neutral, and advantageous. Deleterious mutations decrease the fitness of the host, which may cause their quick elimination in the environment (Loewe & Hill, 2010). Meanwhile, neutral mutations are not affected by selections because of their small effect, and advantageous mutations increase the fitness of the host, allowing them a higher chance of survival (Eyre-Walker & Keightley, 2007; Loewe & Hill, 2010). The resulting mutants remain to strive in the environment depending on the intensity of genetic drift and purifying selection (Webster et al., 2023; Kimura et al., 1963). Thus, variations in the mitochondrial genome of BPH allow the development of diverse phenotypes that can confer an advantage to a specific environment, enabling the species to better adapt in response to the changing environment.



Figure 3. Multiple amino acid sequence alignment of the mitochondrial DNA cytochrome oxidase I gene of the brown planthopper, *Nilaparvata lugens* (Stål) collected from Bohol, Cebu, and Negros Oriental, Visayas, Philippines.

Understanding and considering the genetic diversity in pest populations is important in creating an effective and more efficient pest management strategy. The occurrence of biotypes in BPH is defined by the differences in their non-morphological traits as an effect of the variations in their genetic makeup (Jena & Kim, 2010). Different BPH biotypes show variations in virulence (Claridge & Den Hollander, 1980), resistance development (Myint et al., 2009), and host

preference for feeding or oviposition (Jena & Kim, 2010). Hence, the existing local BPH management could be affected, and perhaps some management strategies could not be effective due to the occurrence of biotypes that already adapted to the management strategies in that locality.

The pairwise fixation index (F_{st}) values, which measure the population differentiation of the BPH population were determined (Table 2). The results showed that F_{st} values between populations Cebu and Bohol ($F_{st}=0.14931$, $P=0.32324$) and Cebu and Negros Oriental ($F_{st}=0.10087$, $P=0.30078$) had shown moderate genetic differentiation, while Bohol and Negros Oriental ($F_{st}=0.04454$, $P=0.42969$) population had shown little genetic differentiation. However, all the compared populations were found to be non-significant. The non-significance of the F_{st} values may indicate that there is no genetic differentiation between the BPH populations. However, due to the small number of samples used in the study, it is suggested to increase the sample size and conduct a wider sample collection.

Table 2. Pairwise genetic p -distances (upper right) and pairwise F_{st} (lower left) among populations of brown planthopper, *Nilaparvata lugens* (Stål) collected from Bohol, Cebu, and Negros Oriental, Visayas, Philippines.

PROVINCES	BOHOL	CEBU	NEGROS ORIENTAL
Bohol		0.32324	0.42969
Cebu	0.14931		0.30078
Negros Oriental	0.04454	0.10087	

Levels of genetic differentiation of F_{st} values: between 0 to 0.05 – little differentiation; between 0.05 to 0.15 – moderate differentiation; between 0.15 to 0.25 – great differentiation; 1 – complete isolation; $P < 0.05$.

The AMOVA is a statistical method in population genetics that quantifies the molecular variation within species (Huang et al., 2021). Table 3 shows the AMOVA of the three different populations of BPH, wherein the Tamura-Nei model was used in the molecular distance with 1000 permutations for the level of significance. The results showed that the contribution of genetic variation within the BPH population was 91.32%, which means great variation is accounted to the difference of BPH within the three localities. On the other hand, the contribution of genetic variation among populations was 8.68%. Therefore, *COI*-based genetic differentiation of the BPH populations studied originated from intra-populations. The genetic differentiation (0.08680), which was measured by the fixation index (F_{st}) as calculated by Arlequin (Excoffier et al., 2005), showed a moderate degree of genetic differentiation among BPH samples from the three localities.

Table 3. Analysis of molecular variance of the brown planthopper, *Nilaparvata lugens* (Stål), collected from Bohol, Cebu, and Negros Oriental, Visayas, Philippines.

SOURCE OF VARIATION	D.F.	SUM OF SQUARES	VARIANCE OF COMPONENTS	PERCENTAGE OF VARIATION
Among populations	2	2.607	0.08595 Va	8.68
Within populations	11	9.947	0.90431 Vb	91.32
Total	13	12.554	0.99026	

Significance tests based on 1023 permutations ($p > 0.05$)
 Fixation Index $F_{st} = 0.08680$

Analysis of the *COI* nucleotide sequence of the fourteen individuals from the three provinces in Central Visayas, Philippines revealed a total of five distinct haplotypes (Table 1 and Figure 4). Haplotype 1 was the most common haplotype shared by two populations (Cebu and Bohol), representing 42.86% of the total *COI* gene sequenced. The second most abundant was Haplotype 3, shared by Negros Oriental and Cebu with 21.43% of the total *COI* gene sequenced. Haplotype 2, with 14.29% of the total *COI* gene sequenced, is unique to Bohol. Haplotype 5, with 14.29% of the total *COI* gene sequenced, is unique to Negros Oriental. Haplotype 4 was considered a singleton haplotype found in Negros Oriental.

The *COI* gene sequences of the five haplotypes were compared with the reported *COI* gene sequences in the GenBank, and all the samples were identified as *N. lugens*, as their nucleotide sequences were >99% to 100% identical to *N. lugens* with 100% query cover. The unrooted network of the five *COI* haplotypes found in three different provinces in Visayas, Philippines, is shown in Figure 5. The size of each circle is proportional to the number of samples with the observed haplotype, while each color represents the portion of samples collected from a specific location. The lines present across the branches represent the nucleotide that differs between the haplotypes. Haplotype 1 differs from haplotypes 4 and 3 by 1 nucleotide and 2 nucleotides, respectively. While Haplotypes 2 and 5, which were not directly connected to Haplotype 1, differed in two to three nucleotides. The major haplotype of BPH in Central Visayas is Haplotype 1, which is considered to be the original or ancestral haplotype in the region (Guru-Pirasanna-Pandi, 2022).




Figure 4. Geographic distribution of the five mitochondrial DNA *cytochrome oxidase I* haplotypes observed from brown planthopper, *Nilaparvata lugens* (Stål) collected in rice fields in Cebu, Bohol, Negros Oriental, Visayas, Philippines.




Figure 5. The haplotype network of the brown planthopper, *Nilaparvata lugens* (Stål), using the sequences of mitochondrial *cytochrome oxidase I*.

Evolutionary changes in the genetic makeup of the insect are induced by several selective pressures in its environment. This includes abiotic and biotic factors that affect the survivability of the insect. If the induced changes bring functional attributes to the pests, the variation can be distributed to insect populations (Karthika et al., 2017). Thus, the presence of the different haplotypes in the three populations may be the result of the insect's adaptation to the environment due to differences in control management in BPH.

Pathogenicity of *B. bassiana* to *N. lugens*

B. bassiana with a fungal concentration of 1×10^8 conidia/mL caused successful infection against BPH populations. The progression of the fungal infection in the three different populations of BPH is shown in Figure 6. The infection of *B. bassiana* to BPH populations started at 24 hrs after treatment with an increasing trend of disease progression of up to 60% mortality at 72 hrs after treatment. The mortality of all the set-ups treated with *B. bassiana* was higher than 50% after 72 hours of post-infection. Thus, this finding aligns with the study of Atta et al. (2020) wherein *B. bassiana* at 1×10^8 conidia/mL sprayed to rice stem pieces caused more than 50% mortality of BPH.




Figure 6. Cumulative corrected percent mortality in Bohol, Cebu, and Negros Oriental populations of adult brown planthopper, *Nilaparvata lugens* (Stål) after exposure to *Beauveria bassiana* (Bals.-Criv) Vuill. with a conidial concentration of 1×10^8 conidia/mL.

Upon performing one-way ANOVA, there were no significant differences in the mortality of the treated BPH between populations ($p = 0.9785$; $F = 0.02172$). Table 4 summarizes the mean lethal time due to exposure of BPH to *B. bassiana*. The values calculated range from 4.08 to 4.26 days, with no significant difference in the mean lethal time of the BPH challenged with *B. bassiana*. This finding suggests that BPH populations were equally susceptible to *B. bassiana*. In addition, the mycosis was confirmed in BPH cadavers by observing the white fungal growth and a stiff body (Figure 7). The stiffness of the insect body is due to the fluid adsorption of the fungus (Skinner et al., 2014).

Table 4. Calculated mean lethal time values of the laboratory bioassays of *Beauveria bassiana* (Bals.-Criv) against the three different populations of *Nilaparvata lugens* (Stål) collected from Bohol, Cebu, and Negros Oriental, Visayas, Philippines.

CALCULATED MEAN LETHAL TIME (DAYS \pm SE)		
Bohol	Cebu	Negros Oriental
4.17 \pm 0.19	4.08 \pm 0.20	4.26 \pm 0.19




Figure 7. The cadavers of adult brown planthopper, *Nilaparvata lugens* (Stål), showing mycosis at ventral (a) and dorsolateral (b) sides after successful infection of *Beauveria bassiana* (Bals.-Criv.) at a conidial concentration of 1×10^8 conidia/mL.

The results showed that *B. bassiana* isolated from a rice bug, *L. oratorius* (Fabricus), was able to cross-infect adult BPH. This conforms with the study of Uma Devi et al. (2008), which reported that *B. bassiana* is a generalist and can be used as a wide-ranged mycoinsecticide after infecting nine different insect hosts. In the Philippines, *B. bassiana* were able to infect fall armyworm, *Spodoptera frugiperda* (Noctuidae), and caused lethal infection in the insect's

larval instars ranging from 23.64% to 97.42% (Montecalvo and Navasero, 2021). The results of the study with regards to the lethal effect of the *B. bassiana* isolated from *L. oratorius* infecting *N. lugens* shows the promising potential of the fungus as a biological control agent for BPH populations in Visayas, Philippines. *B. bassiana* could be incorporated into integrated pest management due to its several advantages. Using EPF is beneficial as they target specific pests and reduce the risk of insecticide resistance development.

The BPH populations with high genetic diversity were hypothesized to exhibit lower mortality when subjected to *B. bassiana*. Insect populations with higher levels of genetic diversity are generally more capable of adapting to environmental changes than those with lower genetic diversity (Birader, 2023) due to the insects' behavioral adaptations and immune capacity (Zhang et al., 2019). However, our findings have shown no significant differences in mortality of the three different BPH populations challenged with *B. bassiana* even the occurrence of genetic variations. Perhaps the occurrence of moderate genetic differentiation among the BPH populations does not affect their susceptibility to *B. bassiana*. Although some studies have shown that immunity of an insect population is reflected in its population genetics, for instance, in the study of Whitehorn et al. (2011), which showed the relationship between the genetic diversity of the moss carder bee (*Bombus muscorum*) and its immunity towards parasites. It was found that *B. muscorum* populations with low genetic diversity had a high occurrence of the insect's gut parasite. Similarly, Ebert et al. (2007) observed that the low genetic diversity of *Daphnia magna* increases the probability of parasitic infection. However, Ekroth et al. (2019) concluded that the disparity in successful pathogen infection between high genetic diversity and low genetic diversity populations was more pronounced in field investigations than in laboratory studies. Hence, field experiments should be conducted to verify the virulence of *B. bassiana* and the possible varying susceptibility of the BPH populations in Central Visayas, Philippines, to this beneficial fungus.

ACKNOWLEDGEMENTS

The authors would like to thank the Department of Agriculture-Bureau of Agricultural Research for the financial support; Dr. Barbara L. Caoili for providing insect samples; Mr. Ariel L. Hirao, Mr. Reynaldo Arapan, Ms. Rochelle A. Quiñones, Ms. Samantha Izabelle R. Alforja, Ms. Daniella Jean E. Pamulaklakin, Mr. Albert D. Sanfuego, Mr. Adrian M. Avila, and Ms. Sylvia M. Calderon for the technical assistance during the study.

REFERENCES CITED

- ABBOTT WS. 1925. A method of computing the effectiveness of an insecticide. *Journal of Economic Entomology* 18: 265–267.
- ATTA B, RIZWAN M, SABIR AM, GOGI MD, FAROOQ MA, BATTYA YA. 2020. Efficacy of entomopathogenic fungi against brown planthopper *Nilaparvata lugens* (Stål) (Homoptera: Delphacidae) under controlled conditions. *Gesunde Pflanzen* 72(2): 101-112. doi: 10.1007/s10343-019-00490-6.
- BACKUS EA, SERRANO MS, RANGER CM. 2005. Mechanisms of hopperburn: an overview of insect taxonomy, behavior, and physiology. *Annual Review of Entomology* 50(11):3087-3113. doi:10.1146/annurev.ento.49.061802.123310.
- BALLARD JWO, MELVIN RG. 2010. Linking the mitochondrial genotype to the organismal phenotype. *Molecular Ecology* 19(8): 1523-1539. doi: 10.1111/j.1365-294X.2010.04594.x.
- BIRADER K. 2023. Genetic diversity and the adaptation of species to changing environments. *Journal of Biodiversity and Endangered Species* 11:3.
- BOTTRELL DG, SCHOENLY KG. 2012. Resurrecting the ghost of green revolutions past: the brown planthopper as a recurring threat to high-yielding rice production in tropical Asia. *Journal of Asia-Pacific Entomology* 15(1): 122-140. doi: 10.1016/j.aspen.2011.09.004.
- CABAUATAN PQ, CABUNAGAN RC, CHOI IR. 2009. Rice viruses transmitted by the brown planthopper *Nilaparvata lugens* Stål. In: K.L. Heong and B. Hardy, editors. *Planthoppers: new threats to the sustainability to of intensive rice production systems in Asia*. International Rice Research Institute. p. 357-368.
- CAMUS MF, WOLF JBW, MORROW EH, DOWLING DK. 2015. Single Nucleotides in the mtDNA Sequence Modify Mitochondrial Molecular Function and Are Associated with Sex-Specific Effects on Fertility and Aging. *Current Biology* 25(20): 2717-2722. doi: 10.1016/j.cub.2015.09.012.
- CARLSON CS, THOMAS DJ, EBERLE MA, SWANSON JE, LIVINGSTON RJ, RIEDER MJ, NICKERSON DA. 2005. Genomic regions exhibiting positive

- selection identified from dense genotype data. *Genome Research* 15(11): 1553-1565. doi: 10.1101/gr.4326505.
- CLEMENT M, POSADA D, CRANDALL K. 2000. TCS: A Computer Program to Estimate Gene Genealogies. *Molecular Ecology* 9: 1657-1660. doi: 10.1046/j.1365-294x.2000.01020.x.
- DANNON HF, DANNON AE, DOURO-KPINDOU OK, ZINSOU AV, HOUNDETE AT, TOFFA-MEHINTO J, ELEGBEDE IATM, OLOU BD, TAMÒ M. 2020. Toward the efficient use of *Beauveria bassiana* in integrated cotton insect pest management. *Journal of Cotton Research* 3: 1-21. doi: 10.1186/s42397-020-00061-5.
- DE FARIA MR, WRAIGHT SP. 2007. Mycoinsecticides and Mycoracaricides: A comprehensive list with worldwide coverage and internal classification of formulation types. *Biological Control* 43(3): 237-256. doi: 10.1016/j.biocontrol.2007.08.001.
- DEAGLE BE, JARMAN SN, COISSAC E, POMPANON F, TABERLET P. 2014. DNA metabarcoding and the *cytochrome c oxidase subunit I* marker: Not a perfect match. *Biology Letters* 10(9): 1-4. doi: 10.1098/rsbl.2014.0562.
- EBERT D, ALTERMATT F, LASS S. 2007. A short-term benefit for outcrossing in a *Daphnia* metapopulation in relation to parasitism. *Journal of the Royal Society Interface* 4(16): 777-785. doi: 10.1098/rsif.2007.0232.
- EKROTH AKE, RAFALUK-MOHR C, KING KC. 2019. Host genetic diversity limits parasite success beyond agricultural systems: A meta-analysis. *Proceedings of the Royal Society B: Biological Sciences* 286(1911): 1-9. doi: 10.1098/rspb.2019.1811.
- EXCOFFIER LG, LAVAL, SCHNEIDER S. 2005. Arlequin ver. 3.0: An integrated software package for population genetics data analysis. *Evolutionary Bioinformatics Online* 1:47-50.
- EYRE-WALKER A, KEIGHTLEY PD. 2007. The distribution of fitness effects of new mutations. *Nature Reviews Genetics* 8: 610-618. doi: 10.1038/nrg2146.
- GASTEIGER E, GATTIKER A, HOOGLAND C, IVANY I, APPEL RD, BAIROCH A. 2003. ExPASy: The proteomics server for in-depth protein knowledge and analysis. *Nucleic Acids Research* 31(13): 3784-3788. doi: 10.1093/nar/gkg563.

- GHAFFAR MB, PRITCHARD J, FORD-LLOYD B. 2011. Brown planthopper (*N. lugens* Stål) feeding behaviour on rice germplasm as an indicator of resistance. PLoS ONE 6(7). doi: 10.1371/journal.pone.0022137.
- GURU-PIRASANNA-PANDI G, ANANT AK, CHOUDHARY JS, BABU SB, BASANA-GOWDA G, ANNAMALAI M, PATIL N, ADAK T, PANNEERSELVAM P, RATH PC. 2022. Molecular diversity of *Nilaparvata lugens* (Stål.) (Hemiptera: Delphacidae) from India based on internal transcribed spacer 1 gene. Current Science. 122(12): 1392-1400. doi: 10.18520/cs/v122/i12/1392-1400.
- HALL TA. 1999. BioEdit: A user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. Nucleic Acids Symposium Series 41: 95-98.
- HEONG KL. 2009. Are planthopper problems caused by a breakdown in ecosystem services? In: K.L.Heong, B.Hardy, editors, Planthoppers: New Threats to the Sustainability of Intensive Rice Production Systems in Asia, p 221–232.
- HEREWARD JP, CAI X, MATIAS AMA, WALTER GH, XU C, WANG Y. 2020. Migration dynamics of an important rice pest: The brown planthopper (*Nilaparvata lugens*) across Asia—insights from population genomics. Evolutionary Applications 13(9): 2449–2459. doi: 10.1111/eva.13047.
- HUANG K, WANG T, DUNN DW, ZHANG P, SUN H, BAOGUO LI. 2021. A generalized framework for AMOVA with multiple hierarchies and ploidies. Integrative Zoology. 16(1): 33-52. doi: 10.1111/1749-4877.12460.
- JENA KK, KIM SM. 2010. Current status of brown planthopper (BPH) resistance and genetics. Rice. 3: 161–171. doi: 10.1007/s12284-010-9050-y.
- KARTHIKA P, VADIVALAGAN C, KRISHNAVENI N, MURUGAN K, NICOLETTI M, CANALE A, BENELLI G. 2017. Contrasting genetic diversity and intra population polymorphism of the invasive pest *Henosepilachna vigintioctopunctata* (Coleoptera, Coccinellidae): A DNA barcoding approach. Journal of Asia-Pacific Entomology 20(1): 23–29. doi:10.1016/j.aspen.2016.11.011.
- KHOA DB, THANG BX, VAN LIEM N, HOLST N, KRISTENSEN M. 2019. Variation in susceptibility of eight insecticides in the brown planthopper *Nilaparvata lugens* in three regions of Vietnam 2015-2017. PLOS ONE. 13(10):1-14. doi: 10.1371/journal.pone.0204962.

- KIMURA M, MARUYAMA T, CROW JF. 1963. The mutation load in small populations. *Genetics* 48:1303–12.
- LACEY LA, BROOKS WM. 1997. Initial handling and diagnosis of diseased insects. L.A. Lacey, editor, *Manual of Techniques in Insect Pathology*. p 1–15 Academic Press, London.
- LAQUINTA JF, CAOILI BL, ARDEZ KP, GUERRERO MS, NAVASERO MV. 2019. Preliminary findings of molecular variation in brown planthopper, *Nilaparvata lugens* (Stål), among populations from Isabela and Laguna [Philippines]. *Philippine Entomologist* 33(2): 190-191.
- LOEWE L, HILL WG. 2010. The population genetics of mutations: good, bad and indifferent. *Philosophical Transactions of the Royal Society B: Biological Sciences* 365(1544): 1153–1167. doi: 10.1098/rstb.2009.0317.
- LUNT DH, ZHANG DX, SZYMURA JM, HEWLTT M, HEWTTT GM. 1996. The insect *cytochrome oxidase I* gene: evolutionary patterns and conserved primers for phylogenetic studies. *Insect Molecular Biology*. 5(3): 153-165. doi: 10.1111/j.1365-2583.1996.tb00049.x.
- MAMIIT RJ, YANAGIDA J, MIURA T. 2021. Productivity hot spots and cold spots: setting geographic priorities for achieving food production targets. *Frontiers in Sustainable Food Systems* 5: 1-15. doi: 10.3389/fsufs.2021.727484.
- MANTZOUKAS, S., KITSIOU, F., NATSIOPOULOS, D., & ELIOPOULOS, P. A. 2022. Entomopathogenic fungi: interactions and applications. *Encyclopedia*, 2(2): 646–656. doi.org/10.3390/encyclopedia2020044.
- MASCARIN GM, JARONSKI ST. 2016. The production and uses of *Beauveria bassiana* as a microbial insecticide. *World Journal of Microbiology and Biotechnology* 32(11): 177. doi: 10.1007/s11274-016-2131-3.
- MASHHOOR K, LAZAR KV, SHANAS S, RAMESH N. 2015. Genetic structure and phylogenetic status of rice brown plant hopper (BPH), *Nilaparvata lugens* Isolated from Kerala, India. *Journal of Applied Biology & Biotechnology* 3(5): 15-18. doi: 10.7324/jabb.2015.3503.
- MATSUMOTO Y, MATSUMURA M, SATO Y, NODA H. 2012. Mitochondrial sequences of *Nilaparvata lugens* (Stål) and *Sogatella furcifera* (Horvath) (Hemiptera, Delphacidae): differentiation of East and Southeast Asian populations. GenBank: AB572313.1.

- MATSUMOTO Y, MATSUMURA M, SANADA-MORIMURA S, HIRAI Y, SATO Y, NODA H. 2013. Mitochondrial *cox* sequences of *Nilaparvata lugens* and *Sogatella furcifera* (Hemiptera, Delphacidae): Low specificity among Asian planthopper populations. *Bulletin of Entomological Research* 103(4): 382–392. doi: 10.1017/S000748531200082X.
- MIN S, LEE SW, CHOI B, LEE SH, KWON DH. 2014. Insecticide resistance monitoring and correlation analysis to select appropriate insecticides against *Nilaparvata lugens* (Stål), a migratory pest in Korea. *Journal of Asia-Pacific Entomology* 17(4): 711-716. doi.org/10.1016/j.aspen.2014.07.005.
- MONDAL D., GHOSH A, ROY D, KUMAR A, SHAMURAILATPAM D, BERA S, GHOSH R, BANDOPADHYAY P, MAJUMDER A. 2017. Yield loss assessment of rice (*Oryza Sativa* L.) due to different biotic stresses under system of rice intensification (SRI). *Journal of Entomology and Zoology Studies* 5(4): 1974–1980.
- MONTECALVO MP, NAVASERO MM. 2021. Comparative virulence of *Beauveria bassiana* (Bals.) Vuill. and *Metarhizium anisopliae* (Metchnikoff) Sorokin to *Spodoptera frugiperda* (J.E. Smith) (Lepidoptera: Noctuidae). *The Journal of International Society for Southeast Asian Agricultural Sciences* 27(1):15-26.
- MUN JH, SONG YH, HEONG KL, RODERICK GK. 1999. Genetic variation among Asian populations of rice planthoppers, *Nilaparvata lugens* and *Sogatella furcifera* (Hemiptera: Delphacidae): mitochondrial DNA sequences. *Bulletin of Entomological Research*. 89(3):245-253. doi: 10.1017/S000748539900036X.
- NAEEMULLAH M, NAKAMURA C, TAKEDA M. 2015. Mitochondrial *COI* DNA sequence analysis of *Nilaparvata lugens*. GenBank: LC036644.1.
- NAGATA T, MASUDA T. 1980. Insecticide susceptibility and wing-form ratio of the brown planthopper, *Nilaparvata lugens* (Stål) (Hemiptera: Delphacidae) and the white backed planthopper, *Sogatella furcifera* (Horvath) (Hemiptera: Delphacidae) of Southeast Asia. *Applied Entomology and Zoology* 15(1): 10-19. doi: 10.1303/aez.15.10.
- OWNLEY BH, PEREIRA RM, KLINGEMAN WE, QUIGLEY NB, LECKIE BM. 2004. *Beauveria bassiana*, a dual purpose biological control with activity against insect pests and plant pathogens. In: R.T Lartey and A.J. Caesar, editors,

- Emerging Concepts in Plant Health Management. India: Research Signpost. p 255-269.
- PARK J, KWON W, PARK J, KIM HJ, LEE BC, KIM Y, CHOI NJ. 2019. The complete mitochondrial genome of *Nilaparvata lugens* (Stål, 1854) captured in Korea (Hemiptera: Delphacidae). Mitochondrial DNA B Resour 4(1): 1674-1676. GenBank: MK590088.1.
- PENTINSAARI M, SALMELA H, MUTANEN M, ROSLIN T. 2016. Molecular evolution of a widely adopted taxonomic marker (*COI*) across the animal tree of life. Scientific Reports. doi: 10.1038/srep35275.
- RAHMAN MA, JAHAN SMH. 2006. Effectiveness of different approaches of chemical and biological control against brown plant hopper, *Nilaparvata lugens* (Stål). Journal of Entomology 3(2): 136-142. doi: 10.3923/je.2006.136.142.
- RODERICK GK. 1996. Geographic structure of insect populations: gene flow, phylogeography, and their uses. Annual Review of Entomology 41: 325-352. doi: 10.1146/annurev.en.41.010196.001545.
- ROZAS J, FERRER-MATA A, SANCHEZ-DELBARRIO JC, LIBRADO P, GUIRAO-RICO S, RAMOS-ONSINS SE, SANCHEZ-GRACIA A. 2017. DnaSP 6: DNA sequence polymorphism analysis of large data sets. Molecular Biology and Evolution 34(12): 3299-3302. doi: 10.1093/molbev/msx248.
- SANCHEZ FF. 1980. Insect pest management in rice in the Philippines. Tropical agriculture research series: proceedings of a symposium on tropical agriculture researches 14: 35-38. Japan International Research Center for Agricultural Sciences.
- SHARIF MK, BUTT MS, ANJUM FM, KHAN SH. 2014. Rice bran: a novel functional ingredient. Critical Reviews in Food Science and Nutrition 54(6): 807-816. doi: 10.1080/10408398.2011.608586.
- SIMON C, FRATI F, BECKENBACH A, CRESPI B, LIU H, FLOOK P. 1994. Evolution, weighting, and phylogenetic utility of mitochondrial gene sequences and a compilation of conserved polymerase chain reaction Primers. Annals of the Entomological Society of America 87(6): 651-701. doi: 10.1093/aesa/87.6.651.
- SKINNER M, PARKER BL, KIM JS. 2014. Role of Entomopathogenic Fungi in Integrated Pest Management. In: D.P. Abrol, editor, Integrated Pest

- Management: Current Concepts and Ecological Perspective. p 169-191. doi: 10.1016/B978-0-12-398529-3.00011-7
- THOMPSON JD, HIGGINS DG, GIBSON TJ. 1994. CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. *Nucleic Acids Research* 22(11): 4673-4689. doi: 10.1093/nar/22.22.4673
- TRACY EF. 2014. The promise of biological control for sustainable agriculture: a stakeholder-based analysis. *Journal of Science Policy & Governance*. 5(1).
- UMA DEVI K, PADMAVATHI J, UMA MAHESWARA RAO C, KHAN AAP, MOHAN MC. 2008. A study of host specificity in the entomopathogenic fungus *Beauveria bassiana* (Hypocreales, Clavicipitaceae). *Biocontrol Science and Technology* 18(10): 975-989. doi: 10.1080/09583150802450451.
- USDA. 2023. Rice Sector at a Glance. U.S Department of Agriculture Economic Research Service. <https://www.ers.usda.gov>.
- WEBSTER MT, BEAUREPAIRE A, NEUMANN P, STOLLE E. 2023. Population Genomics for Insect Conservation. *Annual Review of Animal Biosciences* 11:115-140. doi: 10.1146/annurev-animal-122221.
- WHITEHORN PR, TINSLEY MC, BROWN MJF, DARVILL B, GOULSON D. 2011. Genetic diversity, parasite prevalence and immunity in wild bumblebees. *Proceedings of the Royal Society B: Biological Sciences* 278(1709): 1195-1202. doi: 10.1098/rspb.2010.1550.
- WINNIE RM, RAFFIUDIN R, WIDIARTA IN, RAUF A. 2020. The genetic structure of *Nilaparvata lugens* (Stål.) in Java populations. *HAYATI Journal of Biosciences* 27(4): 330-336. doi: 10.4308/hjb.27.4.330.
- ZHANG DW, XIAO ZJ, ZENG BP, LI K, TANG YL. 2019. Insect behavior and physiological adaptation mechanisms under starvation stress. *Frontiers in Physiology* 10:163. doi: 10.3389/fphys.2019.00163.
- ZIMMERMANN G. 2007. Review on safety of the entomopathogenic fungi *Beauveria bassiana* and *Beauveria brongniartii*. *Biocontrol Science and Technology* 17(6): 553-596. doi: 10.1080/09583150701309006.